

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 391 502 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

25.02.2004 Bulletin 2004/09

(51) Int Cl.7: **C12N 1/21**, C12N 15/09,

C12N 9/42

(21) Application number: **02728174.0**

(86) International application number:

PCT/JP2002/005151

(22) Date of filing: **28.05.2002**

(87) International publication number:

WO 2002/097064 (05.12.2002 Gazette 2002/49)

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**

(30) Priority: **29.05.2001 JP 2001160520**

(71) Applicant: **Kao Corporation**

Tokyo 103-8210 (JP)

(72) Inventors:

- **SAWADA, Kazuhisa**
Haga-gun, Tochigi 321-3497 (JP)

• **ENDO, Keiji**

Haga-gun, Tochigi 321-3497 (JP)

• **OZAWA, Tadahiro**

Haga-gun, Tochigi 321-3497 (JP)

• **TOHATA, Masatoshi**

Haga-gun, Tochigi 321-3497 (JP)

• **OZAKI, Katsuya**

Haga-gun, Tochigi 321-3497 (JP)

(74) Representative: **VOSSIUS & PARTNER**

Siebertstrasse 4

81675 München (DE)

(54) **HOST MICROORGANISMS**

(57) A microorganism wherein one or more genes selected from the group of genes participating in sporulation in the middle to late stages of sporulation have been deleted or inactivated; and a process for producing a target product (a protein) by use the microorganism. No spore is formed when this microorganism is employed, thereby enabling production of a target product

(a protein) while decreasing energy loss, production of a by-product and specific production speed to decrease unnecessary consumption of a medium. Moreover, the production period can be prolonged, whereby the target product (the protein) can be produced efficiently.

EP 1 391 502 A1

Description

Technical Field

5 **[0001]** The present invention relates to a host microorganism which is useful for the production of useful proteins or polypeptides, and to a recombinant microorganism.

Background Art

10 **[0002]** Microorganisms have been employed in the industrial production of a broad range of useful substances. For example, microorganisms have been used to produce not only alcoholic beverages and foods such as *miso* (fermented soy paste) and *shoyu* (soy sauce), but also amino acids, organic acids, nucleic acid-related substances, antibiotics, sugars, lipids, proteins, and many other types of substances. Use of such substances can be found in a wide range of fields, encompassing foods, pharmaceuticals, daily necessities such as detergents and cosmetics, and a variety
15 of raw materials for producing items through chemical processes.

[0003] One important issue in the industrial production of useful substances through use of microorganisms is improvement in productivity. Thus, as a measure thereof, cultivation of substance-producing microorganisms has been performed through traditional genetic techniques such as mutation. In particular, thanks to progress in microbial genetics and biotechnology, such cultivation of substance-producing microorganisms is now being carried out more efficiently than ever by use of a genetic technologies such as a genetic engineering technology, giving rise to development
20 of host microorganisms useful for genetic recombination. For example, there has been developed a microorganism strain, which resulted from improvement of a microorganism strain *Bacillus subtilis* Marburg No. 168 that had been acknowledged to be safe and excellent.

[0004] Microorganisms harbor a diversity of genes, so that they can adapt themselves to environmental changes in the natural kingdom. Therefore, substance productivity of microorganisms cannot necessarily be said to be efficient in terms of industrial production of proteins or similar substances employing limited types of production medium.

[0005] Also, concerning certain types of microorganisms, there have been established strains in which genes participating in early stage sporulation are singly deleted or inactivated. However, these strains will not be said to be sufficiently improved in productivity.

30 **[0006]** Accordingly, an object of the present invention is to provide a host microorganism with which production of proteins or polypeptides can be increased through removal of genes which are useless or harmful in the production of proteins or polypeptides from the genome or inactivation of such genes. Another object of the present invention is to provide a recombinant microorganism produced by incorporating, into the above-mentioned host microorganism, a gene which codes for a protein or polypeptide and which is ligated to a transcription initiation regulation region, translation initiation regulation region, or secretion signal region at the downstream end of the region. Yet another object of
35 the present invention is to provide a method for producing a protein or polypeptide by use of the recombinant microorganism.

Disclosure of the Invention

40 **[0007]** The present inventors have extensively searched, among a variety of genes encoded on a microorganism genome, for genes which are useless or function harmfully in the production of useful proteins or polypeptides, and have found that productivity of a protein or polypeptide of interest can be enhanced by deleting from the genome a specific gene participating in sporulation or by inactivating the gene, and then incorporating into the microorganism a
45 gene which encodes a target protein or polypeptide and which has been ligated to a suitable transcriptional initiation region, a translational initiation region, or a secretion signal region, as compared with the productivity attained by the microorganism without such deletion or inactivation.

[0008] Accordingly, the present invention provides a microorganism in which one or more genes selected from the genes which participate in sporulation in the middle to late stages of sporulation have been deleted or inactivated; a recombinant microorganism obtained by incorporating, into the gene-deleted or gene-inactivated microorganism, a
50 gene which encodes a protein or polypeptide and which has been ligated to a transcription initiation regulation region, a translation initiation regulation region, or a secretion signal region at the downstream end of the region; and a method for producing a protein or polypeptide through use of the recombinant microorganism.

55 **Best Mode for Carrying Out the Invention**

[0009] No limitations are imposed on the parental microorganism which is used to construct the microorganism of the present invention, so long as it has a gene which participates in sporulation. Preferably, the parental microorganism

is a spore-forming microorganism. The parental microorganism may be a wild type or a mutant. Specific examples include bacteria belonging to the genus *Bacillus* such as *Bacillus subtilis*, bacteria belonging to the genus *Clostridium*, and yeasts, with bacteria belonging to the genus *Bacillus* being preferred. Among them, *Bacillus subtilis* is particularly preferred, in view that its complete genome information has already been elucidated, that techniques of genetic engineering and genomic engineering have been established, and that bacteria belonging to the *Bacillus subtilis* have an ability to secrete proteins outside the cells.

[0010] Examples of the target protein or polypeptide which is produced by use of the microorganism of the present invention include enzymes which are useful for foodstuffs, drugs, cosmetics, detergents, fiber treatment, drugs for medical tests, etc.; and proteins and polypeptides such as physiological active factors.

[0011] Two hundred and fifty or more genes discretely present on the genome have been identified to take part in sporulation. Among them, a target gene to be deleted or inactivated in the present invention is preferably a gene that promotes sporulation, and examples of such a gene include those encoding a sporulation-stage-specific σ -factor, genes participating in expression of any of the σ -factor genes, and genes participating in activation of any of the σ -factors. In addition, genes which are transcribed by any of the σ -factors to thereby participate in promotion of sporulation are also included within the scope of the present invention. In the early stage of sporulation (stages 0 - I), extracellular enzymes such as proteases and amylases have been known to be produced in increased amounts as compared with the amounts produced in logarithmic growth phases. Therefore, a target gene to be deleted or inactivated is preferably one or more genes which are expressed specifically in the middle to late stages of sporulation to thereby participate in sporulation. Specifically, a target gene is preferably one or more genes involved in the sporulation stage II, III, IV, or V, more preferably stage II or III, particularly preferably stage II. The present inventors have found that these genes are not directly involved in production of proteins of interest and are also not required for growth of the microorganisms in ordinary medium for industrial production.

[0012] Such genes of *Bacillus subtilis* are listed in Tables 1 and 2.

[0013] In the present specification, names, sites, base numbers, and functions of the genes are described on the basis of the *Bacillus subtilis* genome database that has been reported in Nature, 390, 249-256 (1997) and also published by JAFAN: Japan Functional Analysis Network for *Bacillus subtilis* (BSORF DB) (<http://bacillus.genome.ad.jp/>).

Table 1

Gene	Site (kb)	Function
<i>sigE</i>	1,604	Stage II, mother cell-specific σ E factor
<i>sigF</i>	2,443	Stage II, forespore-specific σ F factor
<i>spoIISB</i>	1,328	Stage II and subsequent stages, participating in sporulation
<i>spoIIE</i>	71	Stage II, activating forespore-specific σ F factor
<i>sigG</i>	1,605	Stages III-V, forespore-specific σ G factor
<i>spoIVCB</i> - <i>spoIIIC</i>	2,652-2,701	Stages IV-V, mother cell-specific σ K factor

Table 2

Gene	Site (kb)	Function
<i>spoIIIGA</i>	1,604	Stage II, activating mother cell-specific σ E factor
<i>spoIIAA</i>	2,444	Stage II, participating in activation of forespore-specific σ F factor
<i>spoIVFB</i>	2,855	Stages IV-V, activating mother cell-specific σ K factor
<i>SpoIIR</i>	3,794	Stage II, participating in activation of mother cell-specific σ E factor
<i>SpoIIJ</i>	4,213	Stages III-V, participating in activation of forespore-specific σ G factor

[0014] The following genes are considered to be equivalent to the genes listed in Table 1: Genes having the same functions as those of the *Bacillus subtilis* genes listed in Table 1 or 2, and genes derived from other microorganisms, preferably derived from a bacterium which belongs to the genus *Bacillus*, and exhibiting 70% or more homology, preferably 80% or more, more preferably 90% or more homology, with one of the genes listed in Table 1. These genes are included in the genes which are to be deleted or inactivated according to the present invention. Homology between amino acid sequences is calculated through the Lipman-Pearson method (Science, 227, 1435 (1985)).

[0015] When one or more genes selected from among the genes described above are deleted or inactivated, chemical

energy consumption required for sporulation of the microorganism is reduced, production period of proteins or polypeptides is prolonged, or other benefits are obtained, resulting in improved productivity of the proteins or polypeptides.

[0016] No limitations are imposed on the number of the genes which are deleted or inactivated, so long as at least one gene is deleted or inactivated. The number may be three or more, or five or more. The number is preferably two or three, particularly preferably two.

[0017] In order to construct the microorganism of the present invention, one or more genes in addition to the above genes may be deleted or inactivated. Through such a combination, a greater effect in improvement of the productivity could be expected.

[0018] Deletion or inactivation of a gene can be performed through known methods. Examples include a method in which target genes are sequentially deleted or inactivated, and a method in which one or more arbitrary DNA fragments are deleted or mutated for inactivation and the resultant gene is analyzed and evaluated in terms of the protein productivity by means of a suitable technique.

[0019] A target gene is deleted or inactivated through, for example, a homologous recombination method. Specifically, a DNA fragment containing a target gene is obtained through cloning by use of a suitable plasmid vector. The obtained DNA fragment is mutated by, among other methods, deleting the entire region of the gene or a portion of the target gene region through a routine gene engineering technique while retaining the DNA fragments connected to the respective ends of the target gene; by causing a nonsense mutation in the structural gene through base substitution, frameshift mutation or the like; or by isolating the target gene fragment through cloning or PCR and inserting a DNA fragment into the isolated target gene fragment. Subsequently, the mutated DNA fragment is introduced into a parental microorganism, to thereby cause homologous recombination with the parental microorganism genome at both regions adjacent to the target gene at the respective ends thereof. Thus, the target gene on the genome can be substituted by a DNA fragment in which the target gene has been deleted or inactivated.

[0020] Several methods have been reported for deleting or inactivating a target gene through homologous recombination when a bacteria in *Bacillus subtilis* is employed as a parental microorganism for producing the microorganism of the present invention (e.g., Mol. Gen. Genet., 223, 268 (1990)). The host microorganism of the present invention can be obtained through repetition of such a method.

[0021] Deletion or inactivation of one or more arbitrary DNA fragments can also be performed by obtaining one or more arbitrary DNA fragments from a parental microorganism through cloning and performing homologous recombination by use of the fragments in a manner similar to that described above, or alternatively by radiation of a γ -ray to the parental microorganism.

[0022] The recombinant microorganism of the present invention can be obtained by incorporating a gene encoding a target protein or polypeptide (hereinafter referred to as "a target protein or polypeptide gene") to the thus-obtained microorganism (host microorganism) in which one or more genes selected from the genes participating in sporulation in the middle to late stages of sporulation have been deleted or inactivated.

[0023] No limitations are imposed on the target protein or polypeptide gene. Examples of such genes include industrially usable enzymes such as enzymes for producing detergents, food, fibers, feed, and chemicals, for medical use, and for diagnosis, and physiologically active peptides. The industrially usable enzymes may be classified, on the basis of their functions, into oxidoreductases, transferases, hydrolases, lyases, isomerases, ligases/synthetases, and others. Preferred examples of the target protein or polypeptide gene include genes encoding a hydrolase such as a cellulase, an α -amylase, or a protease. Specific examples include cellulases belonging to the family 5 in Classification of polysaccharide hydrolases (Biochem. J., 280, 309 (1991)). Among them, cellulases derived from a microorganism, particularly cellulases derived from a bacterium belonging to the genus *Bacillus* are illustrated. More specific examples include alkaline cellulases derived from a bacterium belonging to the genus *Bacillus* and having a sequence of SEQ ID NO: 1 or 2, and cellulases having a sequence having 70% or more, preferably 80% or more, more preferably 90% or more homology with the sequence of SEQ ID NO: 1 or 2. The homology between amino acid sequences is determined through the Lipman-Pearson method (Science, 227, 1435 (1985)). Examples of the α -amylases include α -amylases derived from a microorganism, and liquefaction-type amylases derived from a bacterium belonging to the genus *Bacillus* are particularly preferred. Examples of the proteases include serine proteases and metal proteases derived from a microorganism, particularly derived from a bacterium belonging to the genus *Bacillus*.

[0024] A target protein or polypeptide gene is desirably ligated, at the upstream end thereof, to a regulation region participating in transcription or translation of the gene or secretion of the gene product, i.e., a transcriptional initiation regulation region containing a promoter and a transcriptional initiation point, a translation initiation region containing a ribosome binding site and an initiation codon, or a secretion signal peptide region in a suitable form. For example, a target protein or polypeptide gene is desirably ligated to the above regulation regions contained in a cellulase gene derived from a bacterium belonging to the genus *Bacillus* which is described in, for example, Japanese Patent Application Laid-Open (*kokai*) No. 2000-210081 or Hei 4-190793, and the above regulation regions contained in a region adjacent to the cellulase gene at the upstream end of the gene and having a length of 1 kb or less, preferably 0.6 kb or less. Specifically, a target protein or polypeptide gene is desirably ligated to, among others, a sequence of SEQ ID

NO: 1 or 2, or a base sequence having a certain degree of homology with SEQ ID NO: 1 or 2 and having a regulation function similar to that described above.

[0025] The recombinant microorganism of the present invention can be obtained by combining a DNA fragment containing a target protein or polypeptide gene with a suitable plasmid vector and incorporating the recombinant plasmid into a host microorganism cell through a routine transformation method. Alternatively, the recombinant microorganism of the present invention can be obtained by using as the DNA fragment a DNA fragment ligated to a suitable homologous region of a host microorganism gene and incorporating the resultant DNA fragment directly into the host microorganism gene.

[0026] Production of a target protein or polypeptide through use of the recombinant microorganism of the present invention may be performed by inoculating the recombinant microorganism into a medium containing an assimilable source of carbon and nitrogen and the other essential components, culturing the recombinant microorganism through a conventional method, and, after completion of culture, collecting and purifying the target protein or polypeptide.

[0027] As described above, a host microorganism of interest in which a sporulation-related gene has been deleted or inactivated can be produced, and a recombinant microorganism of interest can be produced through use of the host microorganism. In addition, by use of the recombinant microorganism, a useful protein or polypeptide can be produced efficiently. An example case in which α -amylase or cellulase is produced through use of *Bacillus subtilis* will next be specifically described.

[0028] For example, when the *sigF* gene of a bacterium *Bacillus subtilis* (768 bp) encoding an RNA polymerase subunit σ F-factor which expresses in a forespore in stage II or subsequent stages of sporulation is to be deleted, the following procedure may be employed.

[0029] In the first step, a genome gene is extracted from a host microorganism of a *Bacillus subtilis* strain. Using the genome gene as a template, a DNA fragment at the upstream of the initiation codon of the *sigF* gene and a DNA fragment at the downstream of the termination codon of the *sigF* gene are joined by a marker gene such as chloramphenicol resistant gene inserted therebetween through SOE (splicing by overlap extension) - PCR (Gene, 77, 61 (1989)) or other methods.

[0030] In the next step, the host bacterium *Bacillus subtilis* is transformed by use of the thus-obtained DNA fragment through a competent method, and the transformant is isolated on the basis of chloramphenicol resistance or other characteristics, to thereby cause homologous recombination in the upstream and downstream regions of the *sigF* gene to give a transformant in which the *sigF* gene on the genome is substituted by a marker gene such as a chloramphenicol resistant gene or the like.

[0031] Thereafter, into the thus-obtained transformant and the original cell line of *Bacillus subtilis* serving as a control, a plasmid containing a gene encoding α -amylase or cellulase is introduced. The thus-obtained recombinant is incubated under suitable conditions, for example, under shaking in a vegetative medium. The supernatant of the culture solution is measured in terms of α -amylase activity or cellulase activity, and its productivity is compared with that of the original cell line of *Bacillus subtilis*, to thereby confirm that an increased amount of the target product can be obtained by deleting the *sigF* gene. When the culture solution is subjected to an isolation and purification procedure, α -amylase and cellulase can be obtained.

Examples

Example 1

[0032] A genomic DNA was extracted from *Bacillus subtilis* 168. Using the extracted gene as a template, a 1.5-kb DNA fragment (A), on the genome, adjacent to the *sigF* gene (Base No. 2442658 \leftarrow 2443425) at the upstream end thereof and a 1.5-kb DNA fragment (B) adjacent to the *sigF* gene at the downstream end thereof were multiplied. Separately, a 0.9-kb DNA fragment (C) containing a chloramphenicol resistant gene was multiplied using plasmid pC194 as a template. The fragments (A), (B), and (C) were serially ligated in this order, through SOE-PCR, to thereby prepare a 3.9-kb DNA fragment. The *Bacillus subtilis* 168 was transformed using the thus-obtained DNA fragment through a competent method. The transformed *Bacillus subtilis* 168 was cultured on an LB agar medium containing chloramphenicol, and the colonies were isolated as a transformant. The resulting transformant was confirmed through PCR and sequencing to have a genome in which the region containing the *sigF* gene (2442632-2443318) had been deleted and substituted by the chloramphenicol resistant gene. Separately, each of the following regions on the genome was deleted and substituted by the chloramphenicol resistant gene in a manner similar to that described above: a region (1604136-1604976) containing the *sigE* gene (1604166 \rightarrow 1604885), a region (1347781-1348081) containing a substantial part of the *spoIISB* gene (1347913 \leftarrow 1348083), a region (70537-73018) containing a substantial part of the *spoIIIE* gene (70536 \rightarrow 73019), a region (1605083-1605877) containing a substantial part of the *sigG* gene (1605025 \rightarrow 1605807), a region (2652156-2652723) containing the *spoIVCB* gene (2652262 \rightarrow 2652732), or a region (2652156-2701031) containing a region from the *spoIVCB* gene to the *spoIIIC* gene (2652262 \rightarrow 2701023), to thereby

prepare a microorganism in which a gene participating in sporulation is deleted.

[0033] A DNA fragment (3.1 kb) of the alkaline cellulase gene derived from *Bacillus* sp. KSM-S237 (Japanese Patent Application Laid-Open (*kokai*) No. 2000-210081) was introduced into a shuttle vector pHY300PLK at the cleavage point of restriction enzyme *Bam*HI, to thereby prepare a recombinant plasmid pHY-S237. The plasmid was incorporated through the protoplast method into each of the gene-deleted microorganisms prepared in Example 1 and *Bacillus subtilis* 168 serving as a control. The thus-obtained microorganism was incubated overnight under shaking at 37°C in an LB medium (10 mL). The resultant culture solution (0.05 mL) was inoculated to 2 × L-maltose medium (50 mL; 2% trypton, 1% yeast extraction, 1% NaCl, 7.5% maltose, 7.5-ppm manganese sulfate 4-5 hydrate, 15-ppm tetracyclin), followed by incubation for three days under shaking at 30°C. After completion of incubation, the cells were removed from the culture solution through centrifugation, and the alkaline cellulase activity of the supernatant was measured to determine the amount of alkaline cellulase secreted outside the cells during incubation. As a result, as shown in Table 3, all of the microorganisms in which a gene participating in sporulation had been deleted were found to secrete an increased amount of alkaline cellulase as compared with the control microorganism *Bacillus subtilis* 168 (wild type).

Table 3

Deleted gene	Site of the gene (kb)	Amount of alkaline cellulase secreted (relative value)
<i>sigE</i>	1,604	217
<i>sigF</i>	2,443	212
<i>spoIISB</i>	1,328	140
<i>spoIIE</i>	71	216
<i>sigG</i>	1,605	163
<i>spoIVCB</i> - <i>spoIIIC</i>	2,652-2,701	141
<i>spoIVCB</i>	2,652	141
None (wild type)	-	100

Industrial Applicability

[0034] When the microorganism of the present invention is employed, no spores are formed. Therefore, the invention enables production of a target protein or target polypeptide while decreasing energy loss, production of by-products and specific production speed to largely decrease unnecessary consumption of a medium. Moreover, the production period of the protein or polypeptide can be prolonged, whereby the target product can be produced efficiently.

SEQUENCE LISTING

5 <110> KAO CORPORATION

 <120> Host microorganisms

10 <130> KS0660

 <140>

 <141>

15 <150> JP P2001-160520

 <151> 2001-05-29

20 <160> 2

 <170> PatentIn Ver. 2.1

25 <210> 1

 <211> 3150

 <212> DNA

 <213> Bacillus sp. KSM-S237

30 <220>

 <221> CDS

 <222> (573).. (3044)

35 <220>

 <221> sig_peptide

 <222> (573).. (659)

40 <220>

 <221> mat_peptide

 <222> (660).. (3044)

45 <400> 1

 galligccga igcaacaggc ttatatttag aggaaattc tttttaaatt gaalacggaa 60

 taaaatcagg taaacaggtc ctgattttat ttttttagat ttttagaga acigaagatt 120

50 gaaataaaag tagaagacaa aggacataag aaaatlgcat tagttttaa ttagaaaaac 180

 gccittttat aattatttat acctagaacg aaaatactgt ttcgaaagcg gtttactata 240

55 aaacctata ttccggctct tttttaaacc agggggtaaa aattcacctt agtatcttaa 300

111caacaig ctataa1aaa 111glaagac gcaataigca 1c1c11111 1acga1ata1 360
 5 glaagcgg11 aac111g1gc 1ata1gccga 111aggaagg ggggiagat1 gag1caag1a 420
 gla1a1a1at agataac11a 1aag11g11g agaagcagga gagcatc1gg g11ac1caca 480
 10 ag1111111a aaac111a1ac gaaagcac11 1cgg1a11gc 11a1gaat11 agc1a111ga 540
 11caat11ac1 11aaaa1a1 11aggagg1a at a1g a1g 11a aga aag aaa aca 593
 Met Met Leu Arg Lys Lys Thr
 -25
 15 aag cag 11g a11 1c1 1cc a11 c11 a11 11a g11 11a c11 c1a 1c1 11a 641
 Lys Gln Leu Ile Ser Ser Ile Leu Ile Leu Val Leu Leu Leu Ser Leu
 -20 -15 -10
 20 111 ccg gca gci c11 gca gca gaa gga aac act cgt gaa gac aat 111 689
 Phe Pro Ala Ala Leu Ala Ala Glu Gly Asn Thr Arg Glu Asp Asn Phe
 -5 -1 1 5 10
 25 aaa cat 11a 11a ggi aat gac aat g11 aaa cgc cct 1c1 gag gci ggc 737
 Lys His Leu Leu Gln Asn Asp Asn Val Lys Arg Pro Ser Glu Ala Gly
 15 20 25
 30 gca 11a caa 11a caa gaa g1c gal gga caa a1g aca 11a g1a gal caa 785
 Ala Leu Gln Leu Gln Glu Val Asp Gly Gln Met Thr Leu Val Asp Gln
 30 35 40
 35 cat gga gaa aaa a11 caa 11a cgt gga a1g ag1 aca cac gga 11a cag 833
 His Gly Glu Lys Ile Gln Leu Arg Gly Met Ser Thr His Gly Leu Gln
 45 50 55
 40 1gg 111 cct gag a1c 11g aat gal aac gca 1ac aaa gci c11 1c1 aac 881
 Trp Phe Pro Glu Ile Leu Asn Asp Asn Ala Tyr Lys Ala Leu Ser Asn
 60 65 70
 45 gal 1gg gal 1cc aat a1g a11 cgt c11 gci a1g 1a1 g1a ggi gaa aat 929
 Asp Trp Asp Ser Asn Met Ile Arg Leu Ala Met Tyr Val Gly Glu Asn
 75 80 85 90
 50 ggg 1ac gci aca aac cct gag 11a a1c aaa caa aga g1g a11 gal gga 977
 Gly Tyr Ala Thr Asn Pro Glu Leu Ile Lys Gln Arg Val Ile Asp Gly
 95 100 105
 55 a11 gag 11a gcg a11 gaa aat gac a1g 1a1 g11 a11 g11 gac 1gg cat 1025
 Ile Glu Leu Ala Ile Glu Asn Asp Met Tyr Val Ile Val Asp Trp His

EP 1 391 502 A1

	110	115	120	
5	ggt cat gcg cca ggt gat cct aga gat cct gtt tat gca ggt gct aaa Val His Ala Pro Gly Asp Pro Arg Asp Pro Val Tyr Ala Gly Ala Lys 125 130 135	1073		
10	gat ttc ttt aga gaa att gca gct tta tac cct aat aat cca cac att Asp Phe Phe Arg Glu Ile Ala Ala Leu Tyr Pro Asn Asn Pro His Ile 140 145 150	1121		
15	att tat gag tta gcg aat gag ccg agt agt aat aat aat ggt gga gca Ile Tyr Glu Leu Ala Asn Glu Pro Ser Ser Asn Asn Asn Gly Gly Ala 155 160 165 170	1169		
20	ggg att ccg aat aac gaa gaa ggt tgg aaa gcg gla aaa gaa tat gct Gly Ile Pro Asn Asn Glu Glu Gly Trp Lys Ala Val Lys Glu Tyr Ala 175 180 185	1217		
25	gat cca att gla gaa atg tta cgt aaa agc ggt aat gca gat gac aac Asp Pro Ile Val Glu Met Leu Arg Lys Ser Gly Asn Ala Asp Asp Asn 190 195 200	1265		
30	att atc att gtt ggt agt cca aac tgg agt cag cgt ccg gac tta gca Ile Ile Ile Val Gly Ser Pro Asn Trp Ser Gln Arg Pro Asp Leu Ala 205 210 215	1313		
35	gct gat aat cca att gat gat cac cat aca atg tat act gtt cac ttc Ala Asp Asn Pro Ile Asp Asp His His Thr Met Tyr Thr Val His Phe 220 225 230	1361		
40	tac act ggt tca cat gct gct tca act gaa agc tat ccg tct gaa act Tyr Thr Gly Ser His Ala Ala Ser Thr Glu Ser Tyr Pro Ser Glu Thr 235 240 245 250	1409		
45	cct aac tct gaa aga gga aac gla atg agt aac act cgt tat gcg tta Pro Asn Ser Glu Arg Gly Asn Val Met Ser Asn Thr Arg Tyr Ala Leu 255 260 265	1457		
50	gaa aac gga gla gcg gla ttt gca aca gag tgg gga acg agt caa gct Glu Asn Gly Val Ala Val Phe Ala Thr Glu Trp Gly Thr Ser Gln Ala 270 275 280	1505		
55	agt gga gac ggt ggt cct tac ttt gat gaa gca gat gla tgg att gaa Ser Gly Asp Gly Gly Pro Tyr Phe Asp Glu Ala Asp Val Trp Ile Glu 285 290 295	1553		

EP 1 391 502 A1

5	<p> <i>ttt tta aat gaa aac aac att agc tgg gct aac tgg tct tta acg aat</i> 1601 Phe Leu Asn Glu Asn Asn Ile Ser Trp Ala Asn Trp Ser Leu Thr Asn 300 305 310 </p>
10	<p> <i>aaa aat gaa gta tct ggt gca ttt aca cca ttc gag tta ggt aag tct</i> 1649 Lys Asn Glu Val Ser Gly Ala Phe Thr Pro Phe Glu Leu Gly Lys Ser 315 320 325 330 </p>
15	<p> <i>aac gca acc aat ctt gac cca ggt cca gat cat gtg tgg gca cca gaa</i> 1697 Asn Ala Thr Asn Leu Asp Pro Gly Pro Asp His Val Trp Ala Pro Glu 335 340 345 </p>
20	<p> <i>gaa tta agt ctt tct gga gaa tat gta cgt gct cgt att aaa ggt gtg</i> 1745 Glu Leu Ser Leu Ser Gly Glu Tyr Val Arg Ala Arg Ile Lys Gly Val 350 355 360 </p>
25	<p> <i>aac tat gag cca atc gac cgt aca aaa tac acg aaa gta ctt tgg gac</i> 1793 Asn Tyr Glu Pro Ile Asp Arg Thr Lys Tyr Thr Lys Val Leu Trp Asp 365 370 375 </p>
30	<p> <i>ttt aat gat gga acg aag caa gga ttt gga gtg aat tgg gat tct cca</i> 1841 Phe Asn Asp Gly Thr Lys Gln Gly Phe Gly Val Asn Ser Asp Ser Pro 380 385 390 </p>
35	<p> <i>aat aaa gaa ctt att gca gtt gat aat gaa aac aac act ttg aaa gtt</i> 1889 Asn Lys Glu Leu Ile Ala Val Asp Asn Glu Asn Asn Thr Leu Lys Val 395 400 405 410 </p>
40	<p> <i>tgg gga tta gat gta agt aac gat gtt tca gat ggc aac ttc tgg gct</i> 1937 Ser Gly Leu Asp Val Ser Asn Asp Val Ser Asp Gly Asn Phe Trp Ala 415 420 425 </p>
45	<p> <i>aat gct cgt ctt tct gcc aac ggt tgg gga aaa agt gtt gat att tta</i> 1985 Asn Ala Arg Leu Ser Ala Asn Gly Trp Gly Lys Ser Val Asp Ile Leu 430 435 440 </p>
50	<p> <i>ggt gct gag aag ctt aca atg gat gtt att gtt gat gaa cca acg acg</i> 2033 Gly Ala Glu Lys Leu Thr Met Asp Val Ile Val Asp Glu Pro Thr Thr 445 450 455 </p>
55	<p> <i>gta gct att gcg gcg att cca caa agt agt aaa agt gga tgg gca aat</i> 2081 Val Ala Ile Ala Ala Ile Pro Gln Ser Ser Lys Ser Gly Trp Ala Asn 460 465 470 </p>
60	<p> <i>cca gag cgt gct gtt cga gtg aac gcg gaa gat ttt gtc cag caa acg</i> 2129 Pro Glu Arg Ala Val Arg Val Asn Ala Glu Asp Phe Val Gln Gln Thr </p>

EP 1 391 502 A1

	475	480	485	490	
5	gac ggt aag tat aaa gct gga tta aca att aca gga gaa gat gct cct				2177
	Asp Gly Lys Tyr Lys Ala Gly Leu Thr Ile Thr Gly Glu Asp Ala Pro				
		495	500	505	
10	aac cta aaa aat atc gct ttt cat gaa gaa gat aac aat atg aac aac				2225
	Asn Leu Lys Asn Ile Ala Phe His Glu Glu Asp Asn Asn Met Asn Asn				
		510	515	520	
15	atc att cgt ttc ggt gga act gat gca gct gac gtt att tac tta gat				2273
	Ile Ile Leu Phe Val Gly Thr Asp Ala Ala Asp Val Ile Tyr Leu Asp				
		525	530	535	
20	aac att aaa gta att gga aca gaa gtt gaa att cca gtt gtt cat gat				2321
	Asn Ile Lys Val Ile Gly Thr Glu Val Glu Ile Pro Val Val His Asp				
		540	545	550	
25	cca aaa gga gaa gct gtt ctt cct tct gtt ttt gaa gac ggt aca cgt				2369
	Pro Lys Gly Glu Ala Val Leu Pro Ser Val Phe Glu Asp Gly Thr Arg				
		555	560	565	570
30	caa ggt tgg gac tgg gct gga gag tct ggt ggt aaa aca gct tta aca				2417
	Gln Gly Trp Asp Trp Ala Gly Glu Ser Gly Val Lys Thr Ala Leu Thr				
		575	580	585	
35	att gaa gaa gca aac ggt tct aac gcg tta tca tgg gaa ttt gga tat				2465
	Ile Glu Glu Ala Asn Gly Ser Asn Ala Leu Ser Trp Glu Phe Gly Tyr				
		590	595	600	
40	cca gaa gta aaa cct agt gat aac tgg gca aca gct cca cgt tta gat				2513
	Pro Glu Val Lys Pro Ser Asp Asn Trp Ala Thr Ala Pro Arg Leu Asp				
		605	610	615	
45	ttc tgg aaa tct gac ttg gtt cgc ggt gag aat gat tat gta gct ttt				2561
	Phe Trp Lys Ser Asp Leu Val Arg Gly Glu Asn Asp Tyr Val Ala Phe				
		620	625	630	
50	gat ttc tat cta gat cca gtt cgt gca aca gaa ggc gca atg aat atc				2609
	Asp Phe Tyr Leu Asp Pro Val Arg Ala Thr Glu Gly Ala Met Asn Ile				
		635	640	645	650
55	aat tta gta ttc cag cca cct act aac ggg tat tgg gta caa gca cca				2657
	Asn Leu Val Phe Gln Pro Pro Thr Asn Gly Tyr Trp Val Gln Ala Pro				
		655	660	665	

EP 1 391 502 A1

	aaa acg tat acg att aac ttt gat gaa tta gag gaa gcg aat caa gta	2705
	Lys Thr Tyr Thr Ile Asn Phe Asp Glu Leu Glu Glu Ala Asn Gln Val	
5	670 675 680	
	aat ggt tta tat cac tat gaa gig aaa att aac gta aga gat att aca	2753
	Asn Gly Leu Tyr His Tyr Glu Val Lys Ile Asn Val Arg Asp Ile Thr	
10	685 690 695	
	aac att caa gat gac acg tta cta cgt aac atg atg atc att ttt gca	2801
	Asn Ile Gln Asp Asp Thr Leu Leu Arg Asn Met Met Ile Ile Phe Ala	
15	700 705 710	
	gat gta gaa agt gac ttt gca ggg aga gtc ttt gta gat aat gtt cgt	2849
	Asp Val Glu Ser Asp Phe Ala Gly Arg Val Phe Val Asp Asn Val Arg	
20	715 720 725 730	
	ttt gag ggg gct gct act act gag ccg gtt gaa cca gag cca gtt gat	2897
	Phe Glu Gly Ala Ala Thr Thr Glu Pro Val Glu Pro Glu Pro Val Asp	
25	735 740 745	
	ccg ggc gaa gag acg cca cct gtc gat gag aag gaa gcg aaa aaa gaa	2945
	Pro Gly Glu Glu Thr Pro Pro Val Asp Glu Lys Glu Ala Lys Lys Glu	
30	750 755 760	
	caa aaa gaa gca gag aaa gaa gag aaa gaa gca gta aaa gaa gaa aag	2993
	Gln Lys Glu Ala Glu Lys Glu Glu Lys Glu Ala Val Lys Glu Glu Lys	
	765 770 775	
35	aaa gaa gct aaa gaa gaa aag aaa gca gtc aaa aat gag gct aag aaa	3041
	Lys Glu Ala Lys Glu Glu Lys Lys Ala Val Lys Asn Glu Ala Lys Lys	
	780 785 790	
40	aaa taatctatla aactagttat agggtaatct aaaggctga tglagatctt	3094
	Lys	
	795	
45	tlagataacc tttttcttgc ataactggac acagagttgt tattaaagaa agtaag	3150
	<210> 2	
50	<211> 3332	
	<212> DNA	
	<213> Bacillus sp. KSM-64	
	<220>	
55	<221> CDS	

<222> (610)..(3075)

5

<220>

<221> sig_peptide

<222> (610)..(696)

10

<220>

<221> mat_peptide

<222> (697)..(3075)

15

<400> 2

aglacilacc aiiilagagi caaaagalag aagccaagca ggatligccg atgcaaccgg 60

ciiatattia gagggaaatt ciiiiiaaal igaalacgga alaaaaicag glaaacaggt 120

20

cctgaaiiia tiiiiitgaa tiiiiitgag aactaaagat igaalagaa glagaagaca 180

acggacataa gaaaatigia tiagittitaa itatagaaaa cgciiitcia taattattia 240

25

lacctagaac gaaaalacig tiicgaaagc ggitiactat aaaacciat aticcggtc 300

iiiiitaaa cagggggiga aaatcacic tagiaticia atitcaacat gctataataa 360

30

atigtiaaga cgcaataiac atciiiiit italatatit glaagcggt aacctigtc 420

iatatgccga ittaggaagg ggtagattg aglcaagiag tcataattia galaaciat 480

35

aaglitiga gaagcaggag agaalciggg tiacacaaa giiiiitaaa acatitcga 540

aagcaciic ggtaigcct atgaattag clattigatt caattacit aataattia 600

ggaggiaat atg atg tia aga aag aaa aca aag cag itg att tct tcc att 651

40

Met Met Leu Arg Lys Lys Thr Lys Gln Leu Ile Ser Ser Ile

-25

-20

cit att tia gti tia cit cia tct tia itt ccg aca gct cit gca gca 699

45

Leu Ile Leu Val Leu Leu Leu Ser Leu Phe Pro Thr Ala Leu Ala Ala

-15

-10

-5

-1 1

gaa gga aac aci cgt gaa gac aat itt aaa cat tia tia ggt aat gac 747

50

Glu Gly Asn Thr Arg Glu Asp Asn Phe Lys His Leu Leu Gly Asn Asp

5

10

15

aat gti aaa cgc cct tct gag gct ggc gca tia caa tia caa gaa gtc 795

55

Asn Val Lys Arg Pro Ser Glu Ala Gly Ala Leu Gln Leu Gln Glu Val

20

25

30

EP 1 391 502 A1

5	gat gga caa atg aca tta gta gat caa cat gga gaa aaa att caa tta	843
	Asp Gly Gln Met Thr Leu Val Asp Gln His Gly Glu Lys Ile Gln Leu	
	35 40 45	
10	cgt gga atg agt aca cac gga tta caa tgg ttt cct gag atc ttg aat	891
	Arg Gly Met Ser Thr His Gly Leu Gln Trp Phe Pro Glu Ile Leu Asn	
	50 55 60 65	
15	gat aac gca tac aaa gct ctt gct aac gat tgg gaa tca aat atg att	939
	Asp Asn Ala Tyr Lys Ala Leu Ala Asn Asp Trp Glu Ser Asn Met Ile	
	70 75 80	
20	cgt cta gct atg tat gtc ggt gaa aat ggc tat gct tca aat cca gag	987
	Arg Leu Ala Met Tyr Val Gly Glu Asn Gly Tyr Ala Ser Asn Pro Glu	
	85 90 95	
25	tta att aaa agc aga gtc att aaa gga ata gat ctt gct att gaa aat	1035
	Leu Ile Lys Ser Arg Val Ile Lys Gly Ile Asp Leu Ala Ile Glu Asn	
	100 105 110	
30	gac atg tat gtc atc gtt gat tgg cat gta cat gca cct ggt gat cct	1083
	Asp Met Tyr Val Ile Val Asp Trp His Val His Ala Pro Gly Asp Pro	
	115 120 125	
35	aga gat ccc gtt tac gct gga gca gaa gat ttc ttt aga gat att gca	1131
	Arg Asp Pro Val Tyr Ala Gly Ala Glu Asp Phe Phe Arg Asp Ile Ala	
	130 135 140 145	
40	gca tta tat cct aac aat cca cac att att tat gag tta gcg aat gag	1179
	Ala Leu Tyr Pro Asn Asn Pro His Ile Ile Tyr Glu Leu Ala Asn Glu	
	150 155 160	
45	cca agt agt aac aat aat ggt gga gct ggg att cca aat aat gaa gaa	1227
	Pro Ser Ser Asn Asn Asn Gly Gly Ala Gly Ile Pro Asn Asn Glu Glu	
	165 170 175	
50	ggt tgg aat gcg gta aaa gaa tac gct gat cca att gta gaa atg tta	1275
	Gly Trp Asn Ala Val Lys Glu Tyr Ala Asp Pro Ile Val Glu Met Leu	
	180 185 190	
55	cgt gat agc ggg aac gca gat gac aat att atc att gtt ggt agt cca	1323
	Arg Asp Ser Gly Asn Ala Asp Asp Asn Ile Ile Ile Val Gly Ser Pro	
	195 200 205	
55	aac tgg agt cag cgt cct gac tta gca gct gat aat cca att gat gat	1371

EP 1 391 502 A1

5	Asn Trp Ser Gln Arg Pro Asp Leu Ala Ala Asp Asn Pro Ile Asp Asp 210 215 220 225	
10	cac cal aca aig lat aci gii cac ttc tac aci ggi tca cai gci gci His His Thr Met Tyr Thr Val His Phe Tyr Thr Gly Ser His Ala Ala 230 235 240	1419
15	tca aci gaa agc lat ccg cct gaa aci cct aac tct gaa aga gga aac Ser Thr Glu Ser Tyr Pro Pro Glu Thr Pro Asn Ser Glu Arg Gly Asn 245 250 255	1467
20	gia atg agi aac aci cgt lat gcg tta gaa aac gga gia gca gia ttt Val Met Ser Asn Thr Arg Tyr Ala Leu Glu Asn Gly Val Ala Val Phe 260 265 270	1515
25	gca aca gag tgg gga aci agc caa gca aat gga gat ggi ggi cct tac Ala Thr Glu Trp Gly Thr Ser Gln Ala Asn Gly Asp Gly Gly Pro Tyr 275 280 285	1563
30	ttt gat gaa gca gat gia tgg att gag ttt tta aat gaa aac aac att Phe Asp Glu Ala Asp Val Trp Ile Glu Phe Leu Asn Glu Asn Asn Ile 290 295 300 305	1611
35	agc tgg gci aac tgg tct tta acg aat aaa aat gaa gla tct ggi gca Ser Trp Ala Asn Trp Ser Leu Thr Asn Lys Asn Glu Val Ser Gly Ala 310 315 320	1659
40	ttt aca cca ttc gag tta ggt aag tct aac gca aca agt cit gac cca Phe Thr Pro Phe Glu Leu Gly Lys Ser Asn Ala Thr Ser Leu Asp Pro 325 330 335	1707
45	ggg cca gac caa gla tgg gla cca gaa gag tta agt cit tct gga gaa Gly Pro Asp Gln Val Trp Val Pro Glu Glu Leu Ser Leu Ser Gly Glu 340 345 350	1755
50	lat gla cgt gci cgt att aaa ggt gtg aac lat gag cca atc gac cgt Tyr Val Arg Ala Arg Ile Lys Gly Val Asn Tyr Glu Pro Ile Asp Arg 355 360 365	1803
55	aca aaa tac acg aaa gla cit tgg gac ttt aat gat gga acg aag caa Thr Lys Tyr Thr Lys Val Leu Trp Asp Phe Asn Asp Gly Thr Lys Gln 370 375 380 385	1851
	gga ttt gga glg aat gga gat tct cca gti gaa gat gla gti att gag Gly Phe Gly Val Asn Gly Asp Ser Pro Val Glu Asp Val Val Ile Glu 390 395 400	1899

EP 1 391 502 A1

5	aat gaa gcg ggc gct tta aaa ctt tca gga tta gat gca agt aat gat Asn Glu Ala Gly Ala Leu Lys Leu Ser Gly Leu Asp Ala Ser Asn Asp	1947
	405 410 415	
10	gtt tct gaa ggt aat tac tgg gct aat gct cgt ctt tct gcc gac ggt Val Ser Glu Gly Asn Tyr Trp Ala Asn Ala Arg Leu Ser Ala Asp Gly	1995
	420 425 430	
15	tgg gga aaa agt gtt gat att tta ggt gct gaa aaa ctt act atg gat Trp Gly Lys Ser Val Asp Ile Leu Gly Ala Glu Lys Leu Thr Met Asp	2043
	435 440 445	
20	glg att gtt gat gag ccg acc acg gia tca att gct gca att cca caa Val Ile Val Asp Glu Pro Thr Thr Val Ser Ile Ala Ala Ile Pro Gln	2091
	450 455 460 465	
25	ggg cca tca gcc aat tgg gtt aat cca aat cgt gca att aag gtt gag Gly Pro Ser Ala Asn Trp Val Asn Pro Asn Arg Ala Ile Lys Val Glu	2139
	470 475 480	
30	cca act aat ttc gia ccg tta gga gat aag ttt aaa gcg gaa tta act Pro Thr Asn Phe Val Pro Leu Gly Asp Lys Phe Lys Ala Glu Leu Thr	2187
	485 490 495	
35	ala act tca gct gac tct cca tcc tta gaa gct att gcg atg cat gct Ile Thr Ser Ala Asp Ser Pro Ser Leu Glu Ala Ile Ala Met His Ala	2235
	500 505 510	
40	gaa aat aac aac atc aac aac atc att ctt ttt gia gga act gaa ggt Glu Asn Asn Asn Ile Asn Asn Ile Ile Leu Phe Val Gly Thr Glu Gly	2283
	515 520 525	
45	gct gat gtt atc tat tta gat aac att aaa gia att gga aca gaa gtt Ala Asp Val Ile Tyr Leu Asp Asn Ile Lys Val Ile Gly Thr Glu Val	2331
	530 535 540 545	
50	gaa att cca gtt gtt cat gat cca aaa gga gaa gct gtt ctt cct tct Glu Ile Pro Val Val His Asp Pro Lys Gly Glu Ala Val Leu Pro Ser	2379
	550 555 560	
55	gtt ttt gaa gac ggt aca cgt caa ggt tgg gac tgg gct gga gag tct Val Phe Glu Asp Gly Thr Arg Gln Gly Trp Asp Trp Ala Gly Glu Ser	2427
	565 570 575	
55	ggt gtg aaa aca gct tta aca att gaa gaa gca aac ggt tct aac gcg	2475

EP 1 391 502 A1

	Gly Val Lys Thr Ala Leu Thr Ile Glu Glu Ala Asn Gly Ser Asn Ala	
	580 585 590	
5		
	tta tca tgg gaa ttt gga tac cca gaa gla aaa cct agt gal aac tgg	2523
	Leu Ser Trp Glu Phe Gly Tyr Pro Glu Val Lys Pro Ser Asp Asn Trp	
	595 600 605	
10		
	gca aca gct cca cgt tta gat ttc tgg aaa tct gac tlg gtt cgc ggt	2571
	Ala Thr Ala Pro Arg Leu Asp Phe Trp Lys Ser Asp Leu Val Arg Gly	
	610 615 620 625	
15		
	gaa aat gat tat gla act ttt gat ttc tat cta gat cca gtt cgt gca	2619
	Glu Asn Asp Tyr Val Thr Phe Asp Phe Tyr Leu Asp Pro Val Arg Ala	
	630 635 640	
20		
	aca gaa ggc gca atg aat atc aat tta gla ttc cag cca cct act aac	2667
	Thr Glu Gly Ala Met Asn Ile Asn Leu Val Phe Gln Pro Pro Thr Asn	
	645 650 655	
25		
	ggg tat tgg gla caa gca cca aaa acg tat acg att aac ttt gat gaa	2715
	Gly Tyr Trp Val Gln Ala Pro Lys Thr Tyr Thr Ile Asn Phe Asp Glu	
	660 665 670	
30		
	tta gag gaa gcg aat caa gla aat ggt tta tat cac tat gaa gtg aaa	2763
	Leu Glu Glu Ala Asn Gln Val Asn Gly Leu Tyr His Tyr Glu Val Lys	
	675 680 685	
35		
	att aac gla aga gat att aca aac att caa gat gac acg tta cta cgt	2811
	Ile Asn Val Arg Asp Ile Thr Asn Ile Gln Asp Asp Thr Leu Leu Arg	
	690 695 700 705	
40		
	aac atg atg atc att ttt gca gat gla gaa agt gac ttt gca ggg aga	2859
	Asn Met Met Ile Ile Phe Ala Asp Val Glu Ser Asp Phe Ala Gly Arg	
	710 715 720	
45		
	gtc ttt gla gat aat gtt cgt ttt gag ggg gct gct act act gag ccg	2907
	Val Phe Val Asp Asn Val Arg Phe Glu Gly Ala Ala Thr Thr Glu Pro	
	725 730 735	
50		
	glt gaa cca gag cca gtt gat cct ggc gaa gag acg ccg cct gtc gat	2955
	Val Glu Pro Glu Pro Val Asp Pro Gly Glu Glu Thr Pro Pro Val Asp	
	740 745 750	
55		
	gag aag gaa gcg aaa aaa gaa caa aaa gaa gca gag aaa gaa gag aaa	3003
	Glu Lys Glu Ala Lys Lys Glu Gln Lys Glu Ala Glu Lys Glu Glu Lys	
	755 760 765	

gaa gca gla aaa gaa gaa aag aaa gaa gct aaa gaa gaa aag aaa gca 3051
 Glu Ala Val Lys Glu Glu Lys Lys Glu Ala Lys Glu Glu Lys Lys Ala
 770 775 780 785

aic aaa aat gag gct acg aaa aaa taatcctaata aactagttat agggtaatct 3105
 Ile Lys Asn Glu Ala Thr Lys Lys
 790

aaaggctctga tgcagaatctt tttagataacc tttttttgca taactggaca tagaatgggtt 3165

attaaagaaa gcaagggtgtt tatacgaata taaaaaggta gcgaattttaa atgaaacctt 3225

ttaataatgt ctigtatag aatgaatgaag taatttaaga gggggaaacg aagtgaatac 3285

ggaaatttctt agtagaagaa aaacagacca agaaatcttg caagctt 3332

Claims

1. A microorganism in which one or more genes selected from the genes which participate in speculation in the middle to late stages of sporulation have been deleted or inactivated.
2. The microorganism as recited in claim 1, which is a bacterium belonging to the genus *Bacillus*.
3. The microorganism as recited in claim 2, wherein the bacterium belonging to the genus *Bacillus* is *Bacillus subtilis*.
4. The microorganism as recited in any of claims 1 through 3, wherein the genes are expressed in any of stages II, III, IV, and V of sporulation and thus participate in sporulation.
5. The microorganism as recited in any of claims 1 through 4, wherein the gene to be deleted or inactivated is selected from the group consisting of *sigE* gene, *sigF* gene, *spolIE* gene, *spolISB* gene, and *sigG* gene of *Bacillus subtilis*, genes which fall within a region from *spoIVCB* to *spolIIC* of *Bacillus subtilis*, and genes equivalent to any one or more of these genes.
6. A recombinant microorganism obtained by incorporating, into the microorganism as recited in any of claims 1 through 5, a gene which encodes a protein or polypeptide and which has been ligated to a transcription initiation regulation region, a translation initiation regulation region, or a secretion signal region at the downstream end of the region.
7. The recombinant microorganism as recited in claim 6, wherein the transcriptional initiation regulation region, the translational initiation regulation region, or the secretion signal region is derived from a cellulase gene of a bacterium belonging to the genus *Bacillus* or from a 1 kb region extending upstream of the cellulase gene.
8. The recombinant microorganism as recited in claim 6 or 7, wherein the transcription initiation regulation region, the translation initiation regulation region, or the secretion signal region is derived from a cellulase gene having a sequence of SEQ ID NO: 1 or 2 or a sequence having a homology of 70% or more with the sequence of SEQ ID NO: 1 or 2.
9. A method of producing a protein or polypeptide through use of the microorganism as recited in any one of claims 6 through 8.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/05151

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ C12N1/21, C12N15/09, C12N9/42		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁷ C12N1/21, C12N15/09, C12N9/42		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) MEDLINE, BIOSIS(DIALOG), WPI(DIALOG), SwissProt/PIR/GeneSeq/GenBank/EMBL/DDBJ		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	EP 492274 A2 (Eniricerche Societa per Azioni), 13 October, 1992 (13.10.92), & US 6284490 B1 & JP 4-287686 A	<u>1-4</u> 5-9
X Y	KENNEY T.J., Moran C.P. Jr., Organization and regulation of an operon that encodes a sporulation- essential sigma factor in Bacillus subtilis., J.Bacteriol. 1987, Jul.;169(7):3329-39	<u>1-5</u> 6-9
X Y	MIN K.T., Yudkin M.D., Activity of mutant sigma F proteins truncated near the C terminus., J.Bacteriol. 1992 Nov.;174(22):7144-8	<u>1-5</u> 6-9
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not "E" considered to be of particular relevance earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 26 July, 2002 (26.07.02)		Date of mailing of the international search report 13 August, 2002 (13.08.02)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/05151

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	BARAK I., Youngman P., SpoIIE mutants of Bacillus subtilis comprise two distinct phenotypic classes consistent with a dual functional role for the SpoIIE protein., J.Bacteriol. 1996 Aug.;178(16): 4984-9	<u>1-5</u> 6-9
<u>X</u> Y	TAKAMATSU H. et al., The Bacillus subtilis yabG gene is transcribed by SigK RNA polymerase during sporulation, and yabG mutant spores have altered coat protein composition., J.Bacteriol. 2000 Apr.; 182(7):1883-8	<u>1-5</u> 6-9
Y	JP 2000-210081 A (Kao Corp.), 02 August, 2000 (02.08.00), Par. Nos. [0010], [0014]; sequence No.1 (Family: none)	6-9
P,X	KIM J.H. et al., Construction of spore mutants of Bacillus subtilis for the development as a host for foreign protein production., Biotechnology Letters, June 2001, Vol.23, No.12, pages 999 to 1004	1-9

Form PCT/ISA/210 (continuation of second sheet) (July 1998)